

Name: _____

Skittles Decay

You are going to be simulating the radioactive decay of an unstable isotope. Any given atom of that isotope has a 50% chance of decaying over the course of one half-life (the duration of which is a constant for any given isotope; i.e. about 5700 years for ^{14}C , about 700,000,000 years for ^{235}U). For this model, let's assume that a skittles represents an atom. If the side showing the letter "s" is up; the candy represents an undecayed atom of a radioactive isotope. If the blank side is up, it represents a decayed atom (it has become another element). A half-life is a single trial (steps 2-5 below).

Objectives:

- to understand the importance of the law of averages (thus sample size) on the proportion of undecayed-to-decayed atoms
- to observe that the effect of decay in reducing the population size limits the number of half-lives to which the law of averages applies

Procedure:

1. Count out 100 plain skittles that have one side showing the letter "s" and the other blank. Unfit specimens with two "s" sides or two blank sides may be eaten at this point. Set all 100 sample skittles face up and graph the result as trial). This represents atoms of a radioactive isotope in an igneous rock at crystallization.
2. Put the 100 sample skittles into a cup and shake it (don't let them escape).
3. Pour the contents of the cup out onto a white sheet of paper. Count the number of "undecayed" skittles and put those (only those) back into the cup.
4. Record the number of "undecayed" skittles on your group's column of the table and graph it. The "decayed" skittles can be eaten at this point. You have completed a trial.
5. Shake the cup again.
6. Repeat until you have run out of "undecayed" skittles or until you have finished 10 trials.
7. Put the data from your group's column of the table on the board. Record the data from the other groups and calculate the average for each trial.
8. On your graph, connect your group's data with a black line. Graph the class average and connect the data with a blue line. In a line of different color, graph and connect the probable values (given on the table).
9. Discuss the questions at the end of the worksheet with your group, but write down your own answer.


Name: _____

Skittles Decay Lab Report

Class Data Table:

Run	Group 1	Group 2	Group 3	Group 4	Group 5	Group 6	Group 7	Group 8	Class Total	Class Average	Probable Value
0	100	100	100	100	100	100	100	100		100	100
1											50
2											25
3											12.5
4											6.25
5											3.125
6											1.56
7											0.78
8											0.39
9											0.19
10											0.098

Data Graph:



Name: _____

Discussion Questions:

1. Why didn't each group get the same results?
2. What do the probable values represent?
3. Is there anything improbable about the probable values, whether as applies to skittles or to atoms?
4. Which line, the class average or your group's results, most resemble those of the probable values? Why?
5. DO the earlier or later average trials of your class data most resemble the probable values? Why?
6. For how many half-lives do real or simulated atoms behave like probabilistic values? What are the implications for radiocarbon-dating and uranium-dating?
7. Why don't we use radiocarbon dating for dinosaur bones?